


TECHNISCHE UNIVERSITÄT WIEN Vienna University of Technology



Introduction

The VLBI group at the Institute of Geodesy and Geophysics (TU Wien) has developed and released the first version of a new geodetic VLBI data analysis software called VieVS (Vienna VLBI Software). The software allows analysing the VLBI data with highly sophisticated computational approaches and it is arranged in a easy-to-handle userfriendly structure (see poster "Vienna VLBI Software VieVS – Version 1 released" by Plank et al.). One of the next upgrades is the implementation of a new module (vie_glob) for parameter estimation by a so-called global adjustment, where the connection of the single sessions is done by stacking of the normal equations. The time varying parameters (such as Earth orientation or troposphere parameters) are always modelled by piecewise linear offsets at integer fractions of integer hours which simplifies the connection to the next session and easily allows combination in the sense of GGOS. We focus on the determination of time independent geodynamical parameters like the parameters of the solid Earth tides: Love and Shida numbers. Apart from the estimation of the constant nominal values of Love and Shida numbers for the second degree of the tidal potential, VieVS allows to determine frequency dependent values in the diurnal band together with the Free Core Nutation period. In this poster we show first results obtained from the 24-hour IVS R1 and R4 sessions covering 7 years.





Session G2 / EGU2010-4485

Global adjustment of geodetic and geodynamical parameters in the VLBI software VieVS

H. Spicakova, J. Böhm, T. Nilsson, L. Plank, K. Teke, H. Schuh

Institute of Geodesy and Geophysics, TU Vienna, Austria






Roosbeek et al. (1999) or Hinderer et al. (2000).





hana@mars.hg.tuwien.ac.at



cy		Love number h_{21}		
	Tide	IERS values	this paper	Haas and Schuh (1996)
,	Q1	0.6033	0.6033 ± 0.0065	0.560 ± 0.012
)	O_1	0.6026	0.6026 ± 0.0017	0.606 ± 0.002
,	NO_1	0.6004	0.6005 ± 0.0158	0.435 ± 0.030
)	pi_1	0.5882	0.5895 ± 0.0456	0.623 ± 0.090
)	P ₁	0.5823	0.5842 ± 0.0027	0.574 ± 0.005
	K ₁	0.5261	0.5385 ± 0.0049	0.496 ± 0.002
	psi_1	1.0439	1.2023 ± 0.1587	-0.136 ± 0.228
2	phi ₁	0.6623	0.6590 ± 0.0626	0.702 ± 0.121
5	TT ₁	0.6113	0.6104 ± 0.0830	0.934 ± 0.158
-	J_1	0.6105	0.6096 ± 0.0159	0.538 ± 0.031
		•		۲ <u>ــــــــــــــــــــــــــــــــــــ</u>
period		432 <u>sid</u> . days	413 ± 20 <u>sid</u> . days	426.3 ± 20.3 sid. days
$h_{\rm RS}$		-	-0.00217 ± 0.00015	-0.00162 ± 0.00016

- Plank, K. Teke, A. Pany, J. Wresnik, Ichikawa Y Koyama T Gotoh Plans for the Vienna VLBI Software Vie Working Meeting, edited by G. Bourda, P. Charlot, A. Collioud, pp
- Haas, R., H. Schuh: Determination of frequency dependent Love and Shida numbers from VLBI data. Geophys. Res. Lett. Vol. 23 No.
- Iterferometry: studies of the forced nutations of the Earth. J Geophy
- Dehant; Are the free core nutation parameters variable in time?
- McCarthy, D.D., and G. Petit; IERS Conventions 2003, IERS Technical Note 32. Verlag des Bundesamtes fuer Kartographie un
- nutation period stays between 431 and 434 sidereal days. Geophy Res. Letters., 26, 1, pp. 131-134, 1999.
- Wahr, J.M.: Body tides on an elliptical rotating, elastic and oceanles Earth. Geophys. J. R. Astron. Soc., 64, pp. 677-703, 1981.

European Geosciences Union - General Assembly 2010; Vienna, Austria; 02 – 07 May 2010